# New Trends in Diagnosis and Imaging Follow Up of Renal Angiomyolipomas Part I: Classic AMLs

#### **Description**

Angiomyolipomas (AMLs) are composed of three tissue types: vascular, adipose, and muscular, each of which uniquely affects their imaging features and potential impact on clinical outcomes. The authors provide a framework for understanding how the pathological features of AMLs translate into imaging features, which can be used for categorization and for differentiating AMLs from malignant renal cell carcinomas (RCCs), which may have similar appearances.

# **Learning Objectives**

Upon completing this activity, the reader should be able to

- 1. Identify imaging features of AMLs.
- 2. Distinguish AMLs from echogenic RCCs on ultrasound.
- Manage small (<1 cm) echogenic renal masses detected on ultrasound.

#### **Target Audience**

- Radiologists
- Related imaging professionals

#### Authors

John P. McGahan, MD; Charles M. Vollmer, MD Affiliation: Department of Radiology, University of California, Davis, Sacramento, CA

# **Commercial Support**

None

# Accreditation/Designation Statement

This activity has been planned and implemented in accordance with the accreditation requirements and policies of the Accreditation Council for Continuing Medical Education (ACCME) through the joint providership of the Institute for Advanced Medical Education (IAME) and Anderson Publishing. IAME is accredited by the ACCME to provide continuing medical education for physicians. IAME designates this activity for a maximum of .75 AMA PRA Category 1 Credit<sup>™</sup>. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

#### Instructions

- 1. Review this article in its entirety.
- Visit appliedradiology.org/ SAM2.
- 3. Login or create an account.
- Complete the post test and review the discussion and references.

- 5. Complete the evaluation.
- 6. Print your certificate.

# Estimated Time for Completion

1 hour

# **Date of Release and Review**

CME: May 20, 2025

#### **Expiration Date**

CME: May 20, 2026

#### Disclosures

Planner: Erin Simon Schwartz, MD, discloses no relevant financial relationships with any ineligible companies.

Authors: The authors disclose no relevant financial relationships with any ineligible companies.

IAME has assessed conflicts of interest with its faculty, authors, editors, and any individuals who were in a position to control the content of this CME activity. Any relevant financial relationships were mitigated with an independent peer review of this activity, and no conflicts or commercial bias were detected. IAME's planners, content reviewers, and editorial staff disclose no relationships with ineligible entities.

# New Trends in Diagnosis and Imaging Follow-Up of Renal Angiomyolipomas— Part I: Classic AMLs

John P. McGahan, MD, FACR; Anthony F. Chen, MD

#### Introduction

Classification of renal angiomyolipomas (AMLs) may initially seem simplistic. However, there have already been numerous attempts at pathological or imaging-based classification of these tumors, each of which tries to capture the nuances and variation among AML types.

The majority of AMLs are sporadic and asymptomatic and therefore most commonly identified as an incidental finding on imaging. However, AMLs often occur with tuberous sclerosis, a condition where benign tumors may be present in the brain, eye, heart, lungs, and kidneys. While exceedingly rare, AMLs may also be present with pulmonary lymphangioleiomyomatosis (LAM) (Table 1).<sup>1</sup>

The following approach attempts to provide a framework for understanding how the pathological features of AMLs translate to imaging features, which can be used for

**Affiliation:** Department of Radiology, University of California, Davis, Sacramento, California.

**Disclosures:** The authors have no conflicts of interest to disclose. None of the authors received outside funding for the production of this original manuscript and no part of this article has been previously published elsewhere.

categorization and to differentiate AMLs from malignant renal cell carcinomas (RCCs), which may have similar appearances.

#### **AML Tissue Types and Imaging**

Angiomyolipomas are comprised of 3 tissue types: vascular, adipose, and muscular, each of which uniquely affects their imaging features and potential impact on clinical outcomes—for example, the vascular component of the AML may contribute to spontaneous hemorrhage. In rare instances, AMLs may contain benign epithelial cysts or malignant cells (as in the epithelioid type) (Table 2).

The presence of macroscopic fat is a key imaging feature of classic AMLs. While there are fat-poor or fat-invisible AMLs, where macroscopic fat is not present, that present a greater challenge for imaging diagnosis,<sup>1</sup> as the muscular component of these masses contributes significantly to their imaging features (see Part II of this series), they are rare. In Part I of this 2-part series, we will detail some of the classic features of AMLs that contain macroscopic fat.

# Table 1. Etiologies of Renal Angiomyolipomas

	OCCURRENCE
Sporadic	Majority of angiomyolipomas, without associated symptoms
Tuberous sclerosis	Rare
Lymphangioleiomy omatosis	Very rare
Abbreviations: ACR, American College of Radiology: CAR, Canadian Association of	

#### Table 2. Various Types of Renal Angiomyolipomas

Radiologists.

ТҮРЕ	FREQUENCY
Classic	>90% with
	macroscopic fat
Fat-poor	
Hyperattenuating	5%
angiomyolipoma	
Hypoattenuating	Rare
angiomyolipoma	
Fat-invisible	Rare
Angiomyolipoma with	Very rare
epithelial cysts	
Epithelioid	Very rare
angiomyolipoma	
(potentially	
malignant)	

Published: April 1, 2025. https://doi.org/10.37549/AR-D-25-0076

Applied Radiology

© Anderson Publishing, Ltd. All rights reserved. Reproduction in whole or part without express written permission is strictly prohibited

CME

**Figure 1.** Renal cell carcinomas are most commonly round in shape (A), while angiomyolipomas (AMLs) have variable shapes including oval (B), angular interface (C), and "overflowing beer" sign in some peripheral AMLs (D).



**Figure 2.** Renal US in an adult with multiple angiomyolipomas, including 2 that are oval (1, 3) and one that is angular with "overflowing beer" sign (2) (arrow).







#### **US Features**

It is thought that the macroscopic fat in AMLs contributes to their echogenic imaging features on US. Research has shown that echogenic renal masses are usually AMLs or benign entities such as areas of infarction and/or scarring.<sup>2</sup> However, an echogenic renal mass may also be an RCC,<sup>3</sup> and this overlap creates a diagnostic dilemma. Recent studies have demonstrated characteristics that more reliably differentiate AMLs from RCCs on US, including the shape of the mass, whether it has an angular interface, presence of an "overflowing beer" sign, and an echogenic margin (Figure 1).

Sporadic AMLs may be multifocal, with different shapes (Figure 2), compared with sporadic RCCs. For instance, in separate studies comparing endophytic and peripheral AMLs with RCCs, an oval shape was more common with AMLs than with RCCs, which are almost always round (Figure 2).<sup>4,5</sup>

Endophytic AMLs were more likely to have an echogenic margin compared with RCCs. An angular interface was identified with peripheral AMLs and rarely with RCCs (Figure 2). Similarly, an "overflowing beer" sign (Figure 3) was only identified with peripheral AMLs and never with RCCs (Table 3). Both studies showed that over 80% of the patients with sporadic AMLs were female.4,5 AMLs occurring in the context of syndromes such as tuberous sclerosis are usually multiple and have similar shapes to sporadic AMLs.1

Researchers using US radiomics, with software that is not yet commercially available, have shown **Figure 4.** Benefit of contrast-enhanced US. Renal US demonstrating an echogenic interpolar renal mass with an angular interface, corresponding to an angiomyolipoma (AML) (arrow) (A). Renal US showing a more rounded, lower pole renal mass corresponding to a renal cell carcinoma (RCC) (arrows) (B). Contrast-enhanced US showing renal parenchymal enhancement, but little enhancement within the AML (arrow) (C). Contrast-enhanced US showing that the mass enhances as great or greater than the renal parenchyma, with a slightly enhancing rim, indicating an RCC (arrow) (D). Figure is courtesy of Richard G. Barr, MD, PhD, of Northeast Ohio Medical University.



#### **Table 3. US Features to Help Distinguish Angiomyolipomas**

	RENAL CELL CARCINOMA	RENAL Angiomyolipoma
Round	Very common	Present
Oval	Rare	Common
Angular interface	Rare	Present
"Overflowing beer" sign	Almost never	Present
Echogenic margin	Present	Common

# Table 4. CT and MRI Features of Classic (Macroscopic Fat)Angiomyolipomas

СТ	MRI
Fat attenuation < 10 Hounsfield units	T <sub>2</sub> W—hyperintense region due to macroscopic fat
No calcifications	Fat sat—signal drop out due to macroscopic fat
Occasional aneurysms	Chemical shift—India Ink artifact on contrast images due to macroscopic fat/water interface

Figure 5. Coronal contrast-enhanced CT of the abdomen shows a right lower pole mass with solid components and macroscopic fat with Hounsfield units < 10 (arrows), consistent with a large, fat-rich angiomyolipoma.



that some AMLs are more echogenic than RCCs.<sup>6</sup> These authors compared the tumor to cortex echogenicity ratio and 18 other US texture features to differentiate echogenic AMLs from RCCs. They also found that RCCs were larger than AMLs, which may be another distinguishing feature.<sup>6</sup>

Power Doppler US, contrastenhanced US (CEUS), and microvascular flow imaging are evolving technologies that have been shown to be helpful in diagnosing AMLs. For instance, Barr et al showed that CEUS can be useful in distinguishing echogenic renal masses from other masses (Figure 4).<sup>7</sup> Cao et al found CEUS features of fast washout and rim enhancement occurring with RCC had a sensitivity of 95% and a specificity of 91% in distinguishing RCCs from AMLs.<sup>8</sup>

# **Imaging Follow-Up**

There is agreement within much of the medical literature and among most organizations regarding how best to manage incidental echogenic renal masses greater

8

REVIEW

Figure 6. Classic angiomyolipoma (AML) on MRI. Coronal single-shot fast spin echo T2 image of a left renal peripheral AML (curved arrow), largely isointense to retroperitoneal fat (A). Axial T2 fat-suppressed image showing complete loss of signal intensity consistent with macroscopic fat in an AML (curved arrow) of the left kidney (B). Axial in-phase T1 MRI showing high signal intensity AML (curved arrow) in the left kidney (C). Axial out-of-phase MRI showing the "India ink" artifact as a dark line (arrows) at the interface of the macroscopic fat in the AML with the solid-appearing renal tissue (D).



Figure 7. Angular interface on CT/MRI of angiomyolipoma (AML). An elderly patient with a peripheral AML on contrast-enhanced CT (A) and T1 LAVA post-contrast-enhanced MRI (B) showing angular interface (arrows).



than 1 cm in diameter. Thinsection CT should be considered as the next step in imaging evaluation when an echogenic mass is detected, as it can identify



Canadian Association of Radiology (CAR) and the American College

Figure 8. An elderly patient with a simple right renal cyst demonstrating an angular interface (curved arrow). Also note smaller peripheral hypodensity with an angular interface (arrow) probably representing an additional small cyst.



of Radiology (ACR) agree on the recommendations for echogenic masses greater than 1 cm,<sup>10,11</sup> but disagree on smaller echogenic masses. For example, the ACR's statement indicates that masses smaller than 1 cm in diameter are so rarely malignant that they do not require follow-up,<sup>10</sup> whereas the CAR statement recommends annual US for 5 years.11 Given the more recent studies analyzing the morphological features of these common subcentimeter renal masses, then the recommendation of the ACR can be followed if features such as an echogenic margin, oval shape, angular interface, or "overflowing beer" sign are present.

#### **CT/MRI: Classic AMLs**

Most AMLs occur spontaneously in women and contain macroscopic fat. Renal AMLs are a common occurrence in patients with tuberous sclerosis and can appear in patients with LAM. The features of AMLs occurring with either tuberous sclerosis or LAM are similar to those of the various types of spontaneous AMLs.

Fat-rich AMLs are by far the most common type and are definitively diagnosed on CT and MRI. A classic CT feature of the fat-rich AML is CT

Figure 9. An elderly patient with a large retroperitoneal liposarcoma (L) that is poorly encapsulated and displaces the left kidney (arrow) without a capsular defect.



attenuation of -10 Hounsfield units or less within the macroscopical fatty component (Figure 5).<sup>11,12</sup> MRI features of fat-rich AMLs include increased signal intensity on T2-weighted MRI images, but this is not pathognomonic of AMLs. However, utilizing fat-suppression techniques allows the region of high signal intensity to be suppressed on these sequences, confirming the presence of macroscopic fat (Figure 6). Likewise with chemical shift imaging, there will be an "India ink" artifact at the fat/water (renal parenchyma) interface in the same region as the high signal intensity on T2-weighted imaging (Figure 6; Table 4).<sup>13,14</sup>

Morphological features of exophytic renal masses may be useful in distinguishing benign renal masses (cysts and AMLs) from RCCs on MRI.<sup>15</sup> Verma et al showed that the presence of an angular interface of an exophytic mass > 2 cm on T2 MRI was a strong predictor of benignity.<sup>15</sup> This angular **Figure 10.** An elderly patient with a 1-year follow-up contrast-enhanced CT after radiofrequency ablation of a clear cell renal cell carcinoma, showing an avascular mass with surrounding encapsulated fat (arrows).



interface may be identified with AMLs (Figure 7) and with simple cysts (Figure 8), making it a helpful feature in determining the benignity of a mass.

Other distinguishing features include intratumoral calcifications and aneurysms. Intratumoral calcifications are rare within AMLs but may be present with RCCs. Conversely, intratumoral aneurysms may occur with classic AMLs but are rare with RCCs.

Less common renal and/or retroperitoneal masses may also have a macroscopic fatty component. For instance, retroperitoneal liposarcomas, when discovered, are typically large, with an average diameter of 21 cm in one series.<sup>16</sup> They less commonly have an associated renal parenchymal defect than large AMLs, and liposarcomas are more often poorly marginated, sometimes displacing the kidney (Figure 9).<sup>16</sup> Patients treated for RCC by radiofrequency ablation may have encapsulated retroperitoneal fat "pseudomass" at the site of ablation (Figure 10), not to be mistaken for a fatcontaining neoplasm.<sup>17</sup>

#### Conclusion

Understanding the histology and imaging features of classic AMLs can aid in the distinction of these lesions from other primary renal neoplasms, most commonly RCC. While emerging radiomics and advanced US techniques can provide increasingly valuable information, CT and MRI remain the mainstay for imaging characterization of these masses.

#### References

1) Skelton WP, Leslie SW. Renal angiomyolipoma. *StatPearls* [Internet]. StatPearls Publishing. 2024.

2) Kwong A, Mitchell A, Wang D, McGahan JP. Etiology of small echogenic renal masses. *J Ultrasound Med.* 2022;41(10):2567-2575. doi: 10.1002/jum.15946

3) Sidhar K, McGahan JP, Early HM, et al. Renal cell carcinomas: sonographic appearance depending on size and histologic type. *J Ultrasound Med.* 2016;35(2):311-320. doi:10.7863/ultra.15.03051

4) Chen AF, Getz MLD, McGahan JP, Wilson MD, Larson MC. Predictors of benignity for small endophytic echogenic renal masses. *J Ultrasound Med.* 2025;44(3):483-492. doi:10. 1002/jum.16610

5) Chen AF, McGahan JP, Wilson MD, et al. Are there ultrasound features to distinguish small (<3 cm) peripheral renal angiomyolipomas from renal cell carcinomas?. *J Ultrasound Med.* 2023;42(9):2083-2094. doi: 10. 1002/jum.16229

6) Habibollahi P, Sultan LR, Bialo D, et al. Hyperechoic renal masses: differentiation of angiomyolipomas from renal cell carcinomas using tumor size and ultrasound radiomics. *Ultrasound Med Biol.* 2022;48(5):887-894. doi:10.1016/j.ultrasmedbio.2022.01.011

April 2025

7) Barr RG, Peterson C, Hindi A. Evaluation of indeterminate renal masses with contrast-enhanced US: a diagnostic performance study. *Radiology*. 2014;271(1):133-142. doi:10.1148/radiol. 13130161

8) Cao H, Fang L, Chen L, et al. The value of contrast-enhanced ultrasound in diagnosing small renal cell carcinoma subtypes and angiomyolipoma. *J Ultrasound Med.* 2022;41(6):1415-1423. doi:10.1002/jum. 15824

9) SimpsonE, Patel U. Diagnosis of angiomyolipoma using computed tomographyregion of interest < or =-10 HU or 4 adjacent pixels < or =-10 HU are recommended as the diagnostic thresholds. *Clin Radiol.* 2006;61(5):410-416. doi: 10.1016/j. crad.2005.12.013

10) Berland LL, Silverman SG, Gore RM, et al. Managing incidental findings on abdominal CT: white paper of the ACR incidental findings committee. *J Am Coll Radiol*. 2010;7(10):754-773. doi:10.1016/j.jacr.2010.06. 013

11) Kirkpatrick IDC, Brahm GL, Mnatzakanian GN, et al. Recommendations for the management of the incidental renal mass in adults: endorsement and adaptation of the 2017 ACR incidental findings committee white paper by the Canadian association of radiologists incidental findings working group. *Can Assoc Radiol J.* 2019;70(2):125-133. doi:10.1016/j.carj.2019.03.002

12) Park BK. Renal angiomyolipoma: radiologic classification and imaging features according to the amount of fat. *AJR Am J Roentgenol.* 2017;209(4):826-835. doi:10.2214/ AJR.17.17973 13) Jinzaki M, Silverman SG, Akita H, et al. Renal angiomyolipoma: a radiological classification and update on recent developments in diagnosis and management. *Abdom Imaging.* 2014;39(3):588-604. doi:10. 1007/s00261-014-0083-3

14) Thiravit S, Teerasamit W, Thiravit P. The different faces of renal angiomyolipomas on radiologic imaging: a pictorial review. *Br J Radiol.* 2018;91(1084):20170533. doi:10.1259/bjr.20170533

15) Verma SK, Mitchell DG, Yang R, et al. Exophytic renal masses: angular interface with renal parenchyma for distinguishing benign from malignant lesions at MR imaging. *Radiology*. 2010;255(2):501-507. doi: 10.1148/radiol.09091109

16) Woo S, Kim SY, Cho JY, Kim SH, Lee MS. Exophytic renal angiomyolipoma and perirenal liposarcoma: revisiting the role of CT for differential diagnosis. *Acta Radiol.* 2016;57(2):249-255. doi:10.1177/ 0284185115574543

17) Grewal A, Khera SS, McGahan JP, et al. Utility of intraprocedural contrast-enhanced CT in ablation of renal masses. *AJR Am J Roentgenol.* 2020;214(1):122-128. doi:10.2214/ AJR.19.21584